

Honour School of Mathematical and Theoretical Physics Part C  
Master of Science in Mathematical and Theoretical Physics

---

**ADVANCED FLUID DYNAMICS**  
**Trinity Term 2017**

---

**TUESDAY, 18TH APRIL 2017, 2:30pm to 4:30pm**

*You should submit answers to both questions.*

*You must start a new booklet for each question which you attempt. Indicate on the front sheet the numbers of the questions attempted. A booklet with the front sheet completed must be handed in even if no question has been attempted.*

*The numbers in the margin indicate the weight that the Examiners anticipate assigning to each part of the question.*

**Do not turn this page until you are told that you may do so**

1. **Magnetohydrodynamics.** The magnetic field in the solar atmosphere can be thought of as consisting of a family of arc-shaped field lines with footpoints rooted on the surface of the Sun. Below, we will ignore the curvature of the solar surface and denote by  $z$  the spatial coordinate perpendicular to it (the vertical direction).

- (a) [4 marks] Consider a magnetic field of the type described above and argue that a field configuration with a given set of footpoints can be represented as

$$\mathbf{B} = \nabla\alpha \times \nabla\beta, \quad (1)$$

where  $\alpha(x, y, z)$  and  $\beta(x, y, z)$  are some functions whose values at  $z = 0$  are given.

- (b) [9 marks] Show that the field that minimises the magnetic energy within a domain subject to the constraint that  $\alpha$  and  $\beta$  are fixed at the boundary of the domain is a force-free field.
- (c) [3 marks] Show that if this field is a linear force-free field, it satisfies the Helmholtz equation

$$\nabla^2\mathbf{B} + \lambda^2\mathbf{B} = 0, \quad (2)$$

where  $\lambda$  is a constant.

- (d) [9 marks] Consider the semi-infinite domain  $z \geq 0$ . Find a linear force-free field that decays exponentially with  $z$ , is independent of the horizontal coordinate  $y$ , has  $B_z = B_0$  at  $x = z = 0$ , and  $B_x = B_y = 0$  at  $x = 0$  and  $x = a$ . Sketch this field in the  $(x, z)$  and  $(x, y)$  planes. For what values of  $\lambda$  does such a field exist?

## 2. Complex Fluids.

- (a) [5 marks] Consider a suspension of pairs of spheres joined by Hookean springs with spring constant  $H$ . The function  $\psi(\mathbf{x}, \mathbf{R}, t)$  specifies a distribution of sphere-spring pairs with separation  $\mathbf{R}$ , and is normalised so that

$$\int \psi(\mathbf{x}, \mathbf{R}, t) d\mathbf{R} = 1.$$

There are  $n$  such sphere-spring pairs per unit volume in three-dimensional space.

Show that the number of springs per unit area with separation  $\mathbf{R}$  that intersect a plane with normal  $\mathbf{n}$  is

$$n(\mathbf{R} \cdot \mathbf{n})\psi,$$

and hence that the stress due to the stretching of the springs is

$$nH \int \mathbf{R}\mathbf{R} \psi(\mathbf{x}, \mathbf{R}, t) d\mathbf{R} \equiv nH \langle \mathbf{R}\mathbf{R} \rangle.$$

- (b) [13 marks] The total stress due to the sphere-spring pairs is

$$\boldsymbol{\sigma} = -nk_B T \mathbf{I} + nH \langle \mathbf{R}\mathbf{R} \rangle \quad (\dagger)$$

at temperature  $T$ . The effect of the other sphere-spring pairs in the suspension can be modelled by supposing that a sphere moving with velocity  $\mathbf{v}$  relative to its surrounding fluid experiences an anisotropic hydrodynamic resistance force  $-\zeta(\mathbf{I} + \alpha\boldsymbol{\sigma})^{-1}\mathbf{v}$ .

By formulating equations of motion for the two spheres in a sphere-spring pair, subject to Brownian forces, show that  $\psi$  obeys the equation

$$\partial_t \psi + \mathbf{u} \cdot \nabla \psi + \nabla_{\mathbf{R}} \cdot \left( \mathbf{R} \cdot (\nabla \mathbf{u}) \psi - \frac{2}{\zeta} (\mathbf{I} + \alpha\boldsymbol{\sigma}) H \mathbf{R} \psi \right) = \frac{2k_B T}{\zeta} \nabla_{\mathbf{R}} \cdot ((\mathbf{I} + \alpha\boldsymbol{\sigma}) \cdot \nabla_{\mathbf{R}} \psi),$$

where  $[\nabla \mathbf{u}]_{ij} = \partial_i u_j$ , and hence that  $\boldsymbol{\sigma}$  obeys the equation

$$\lambda \left( \partial_t \boldsymbol{\sigma} + \mathbf{u} \cdot \nabla \boldsymbol{\sigma} - \boldsymbol{\sigma} \cdot (\nabla \mathbf{u}) - (\nabla \mathbf{u})^T \cdot \boldsymbol{\sigma} \right) + \boldsymbol{\sigma} + \alpha \boldsymbol{\sigma}^2 = \mu \left( (\nabla \mathbf{u}) + (\nabla \mathbf{u})^T \right). \quad (\star)$$

Give expressions for the coefficients  $\lambda$  and  $\mu$ , and an interpretation of the expression  $(\partial_t \boldsymbol{\sigma} + \dots)$  on the left-hand side.

What property of this equation for  $\boldsymbol{\sigma}$  justifies the coefficient of the isotropic contribution to the stress in  $(\dagger)$ ?

*[You may assume that the Brownian force on the sphere at  $\mathbf{r}_i$  is  $-k_B T \nabla_{\mathbf{r}_i} \psi$ , and neglect any consequent Brownian diffusion in the  $\mathbf{x}$  coordinates.]*

- (c) [7 marks] Suppose the velocity field is  $\mathbf{u} = (\gamma x, -\frac{1}{2}\gamma y, -\frac{1}{2}\gamma z)$ . Show that there are steady, spatially uniform solutions of  $(\star)$  with

$$\boldsymbol{\sigma} = \begin{pmatrix} \sigma_{11} & 0 & 0 \\ 0 & \sigma_{22} & 0 \\ 0 & 0 & \sigma_{22} \end{pmatrix},$$

and find the equations satisfied by  $\sigma_{11}$  and  $\sigma_{22}$ . Comment on the behaviour of  $\sigma_{11}$  as  $\gamma$  increases, for both  $\alpha > 0$  and  $\alpha = 0$ .